

Cold Bose gases with large scattering lengths

S. Cowell¹, H. Heiselberg², I. E. Mazets³, J. Morales¹, V. R. Pandharipande¹, and C. J. Pethick²

¹ *Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Urbana, IL 61801-3080,*

² *NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark,*

³ *Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia.*

We calculate the energy and condensate fraction for a dense system of bosons interacting through an attractive short range interaction with positive s-wave scattering length a . At high densities, $n \gg a^{-3}$, the energy per particle, chemical potential, and square of the sound speed are independent of the scattering length and proportional to $n^{2/3}$, as in Fermi systems. The condensate is quenched at densities $na^3 \simeq 1$.

PACS numbers: 03.75.Fi, 21.65.+f, 74.20.Fg, 67.60.-g

The densities of atomic vapors under experimental conditions are generally low, in the sense that the distance between particles is typically much larger than the magnitude of the atomic scattering length a , which characterizes the strength of two-body interactions. Recently, it has become possible to tune atomic scattering lengths to essentially any value, positive or negative, by exploiting Feshbach resonances [1]. This opens up the possibility of creating systems which are dense, in the sense that the “gaseousness parameter”, $n|a|^3$, where n is the particle density, is large compared with unity [2].

In dilute alkali gases, the bare two-body interaction at distances large compared with the atomic size is attractive due to the van der Waals interaction, but the effective two-body interaction at low energies, which is given by $4\pi\hbar^2 a/m$, may be large and positive due to the existence of a bound state just below threshold.

One of the novel features of such gases is that the characteristic length R associated with the range of the interatomic potential is of order 10^{-6} cm. The quantity $r_0 = (3/4\pi n)^{1/3}$, which is a measure of the atomic separation, is typically of order 10^{-4} cm, and therefore much larger than R . In this Letter we consider the properties of a Bose gas for arbitrary $n|a|^3$, assuming that $R \ll r_0$. The corresponding problem for a Fermi gas has been considered previously [3].

We begin by reviewing briefly the properties of the dilute Bose gas with repulsive interactions, $a > 0$. At low densities ($na^3 \ll 1$) the interaction energy is proportional to the scattering length and the density, as was first found by Lenz [4]. With higher order terms calculated by Huang, Lee, and Yang [5] and Wu [6] included, the energy per particle is

$$\frac{E}{N} = 2\pi \frac{\hbar^2 a}{m} n \left[1 + \frac{128}{15} \left(\frac{na^3}{\pi} \right)^{1/2} + 8 \left(\frac{4\pi}{3} - \sqrt{3} \right) na^3 \ln(na^3) + \mathcal{O}(na^3) \right]. \quad (1)$$

To this order, the only property of the two-body interaction that enters is the scattering length. These results

have been used to calculate the leading corrections beyond mean-field theory to the properties of trapped Bose gases [7]. Terms of order na^3 and higher depend on other properties of the interaction, as discussed in detail recently in Ref. [8].

The expansions leading to the above results are asymptotic, and consequently they give little guidance to the properties of a dense gas, $na^3 \gtrsim 1$. Even though the range of the interaction is small compared with the particle separation, correlations between particles are very important, as one can see from the fact that all terms in the low-density expansion Eq. (1) are formally of the same order of magnitude when $na^3 \simeq 1$.

The Jastrow wave function [10]

$$\Psi_J(\mathbf{r}_1, \dots, \mathbf{r}_N) = \prod_{i < j} f(\mathbf{r}_i - \mathbf{r}_j). \quad (2)$$

provides a good approximation for cold dense Bose systems such as the hard-sphere gas and liquid ^4He . In these systems the pair correlation function $f(r)$ may be determined variationally by minimizing the expectation value of the energy, $E/N = \langle \Psi | H | \Psi \rangle / \langle \Psi | \Psi \rangle$, which may be calculated by Monte Carlo methods [11]. The hard-sphere gas problem has also been solved, essentially exactly, by quantum Monte Carlo methods [12] for scattering lengths almost as large as r_0 .

A much simpler method, the lowest order constrained variational (LOCV) method [13], provides a fair approximation to the results of quantum Monte Carlo calculations. The basic idea of this method is that for $r < r_0$ the Jastrow function $f(r)$ approximately obeys the Schrödinger equation for a pair of particles,

$$\left[-\frac{\hbar^2}{m} \frac{d^2}{dr^2} + v(r) \right] r f(r) = \lambda r f(r), \quad (3)$$

where λ is the pair energy, while for $r \gg r_0$, f tends to unity. To take into account many-body effects, which become important when r is comparable to r_0 , we impose the condition that $f(r > d) = 1$ and $f'(r = d) = 0$ at some healing distance d . In the LOCV method the

expectation value of the energy is approximated by the two-body cluster contribution

$$E/N = 2\pi n \lambda \int_0^d f^2(r) r^2 dr, \quad (4)$$

and d is chosen such that

$$4\pi n \int_0^d f^2(r) r^2 dr = 1. \quad (5)$$

This ensures that on average there are only correlated pairs, and $E/N = \lambda/2$. The energy of a pair is λ , and thus the calculation of the energy by evaluating the expectation value of the Hamiltonian is consistent with the eigenvalue of the Schrödinger equation (3).

When the range of the interaction is small compared to both $|a|$ and d , as in dilute alkali gases, the interaction can be replaced by the boundary condition $(rf)'/rf = -1/a$ at $r = 0$. This boundary condition is also satisfied by the two-body s-wave scattering wave function, and the scattering length a is then the position of the node of $f(r)$ at $\lambda = 0$. The solution to Eq. (3) for positive $\lambda = k^2/m$, is $rf(r) \propto \sin[k(r - b)]$, with a node at $r = b$, and $kb = \arctan ka$. In the dilute limit ($ka \rightarrow 0$), b is equal to a , and in the dense limit ($ka \rightarrow \infty$), $b = \pi/2k$.

The other boundary condition, $f'(r = d) = 0$, gives

$$\frac{a}{d} = \frac{\kappa^{-1} \tan \kappa - 1}{1 + \kappa \tan \kappa}, \quad (6)$$

where $\kappa = kd$, and the value of d is determined from Eq. (5). When $f^2(r) = 1$, Eq. (5) gives $d = r_0$. Since $f(r \sim d) \simeq 1$ the value of d is generally close to r_0 .

In the LOCV approximation $f(r)$ for a short range interaction is therefore given by

$$f(r < d) = \frac{d \sin[k(r - b)]}{r \sin[k(d - b)]}. \quad (7)$$

It is negative for $r < b$, and changes sign at $r = b$. The Jastrow wave function (Eq. 2) does not describe the nodeless ground state in this case. When $a \ll r_0$ the gas has only pairwise collisions, and the Jastrow wave function Ψ_J describes its state accurately. We assume that, as the gas is compressed or a is increased adiabatically, the nodal structure of the wave function is maintained and that Ψ_J continues to be a good approximation to the true wave function. However, as a or n increases, the rate of three-body collisions, which deplete the monoatomic gas density by forming molecules, will increase and limit the lifetime of the monoatomic gas.

The variational principle cannot be used to determine the Jastrow function $f(r)$ in the case of attractive short range interactions, because the gas is not in its ground state. In this case, it may easily be verified that the energy of the gas, as a function of the healing distance d , has an uninteresting minimum at $d \rightarrow 0$ with value zero. Such a minimum exists for all δ -function interactions.

However, we anticipate that for the present problem the LOCV method will provide a good approximation if we take Eq. (7) for the Jastrow factor.

In Fig.1 we show the energy per particle for hard-sphere and attractive short range potentials in the LOCV approximation, along with the low-density expansion (1) (without the logarithmic term) and the Lenz expression as a function of a/r_0 , and the exact quantum Monte Carlo results for hard-spheres. The LOCV energy is close to the low density expansion at small values of a/r_0 , though it does not have the correct analytical form. It contains the leading Lenz term in Eq. (1), as may be verified by expanding the $\tan \kappa$ in Eq. 6 in powers of κ [14], but the $(na^3)^{1/2}$ and higher terms are approximated. The LOCV energies for the hard-sphere gas are in excellent agreement with the exact ones for $a/r_0 \leq 0.6$, but for larger a/r_0 the results for the hard-sphere and the attractive short range potentials deviate significantly.




FIG. 1. Energy per particle, multiplied by $2mr_0^2/\hbar^2$ vs. a/r_0 . The LOCV results for attractive short range and hard-sphere interactions are shown by full and dashed lines respectively, while the dash-dot and dotted lines show the Lenz result and the low density expansion. Monte-Carlo numerical results for a hard-sphere gas are shown as stars.

In the small a/r_0 limit, using the result $b = a$, and $d \sim r_0$ we find that the LOCV correlation function is given by

$$f(r < r_0) = \frac{r_0 \sin[k(r - a)]}{r \sin[k(r_0 - a)]}, \quad (8)$$

with $k^2 = 3a/r_0^3$. At small r this correlation is $\sim 1 - a/r$, as in the two-body scattering state at zero energy.

By contrast, when a/r_0 is large we have $b = \pi/2k$ and Eq. (6) reduces to

$$\kappa \tan \kappa = -1, \quad (9)$$

with solutions $kd = \kappa = 2.798386..., 6.1212..., \dots$, and asymptotically $\kappa = n\pi$ for integer n . The pair correlation function is given by

$$f(r < d) = \frac{d \cos kr}{r \cos kd}. \quad (10)$$

Substituting this expression in Eq. (5) gives $d = (2/3)^{1/3}r_0$ for $kd = 2.798386...$, appropriate for a Jastrow factor f with one node. The energy is given by

$$E/N = \frac{\lambda}{2} = \left(\frac{3}{2}\right)^{2/3} \frac{\hbar^2(kd)^2}{2mr_0^2} = 13.33\hbar^2 \frac{n^{2/3}}{m}. \quad (11)$$

This energy is proportional to $n^{2/3}$ as one would expect on dimensional arguments. A similar result, ($E/N \propto n^{2/3}$), was found for fermions [3] in the $a \rightarrow -\infty$ limit. (We obtain $E/N = 15.26 \hbar^2 n^{2/3}/m$ on approximating d by r_0 .)

For the case of attractive short range interactions it is difficult to go beyond the LOCV calculation. When $a/r_0 < 0.2$ we can perform Monte Carlo calculations of the expectation value of the energy, and the results are not significantly different from those from the LOCV method. At larger values of a/r_0 the state described by the Ψ_J has a clustering instability. All the atoms end up in a few dense clusters as the Metropolis walk progresses, as is expected to happen with time in experiments.

The rate of molecule formation is proportional to the probability of finding three atoms together. In the lowest order (3-body) cluster approximation the probability of finding three atoms at points $\mathbf{r}_1$, $\mathbf{r}_2$, and $\mathbf{r}_3$ is proportional to $f^2(r_{12})f^2(r_{23})f^2(r_{31})$, where $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$. It diverges as all $r_{ij} \rightarrow 0$ for the f given by Eq. (10) or Eq. (8). However, within the interaction radius R the true f does not have the $1/r$ behavior. We can estimate the probability $P_3(R)$ of finding two atoms within the interaction range R of a third by approximating the f at $r < R$ by $f(R)$. In the dense limit Eq. (10) gives $f(r < R) \sim -r_0/R$, and

$$P_3(R) \sim \left(\frac{r_0}{R}\right)^6 \left(\frac{4\pi R^3 n}{3}\right)^2 = 1, \quad (12)$$

independent of the gas density. In contrast, at small values of a/r_0 Eq. (8) gives $f(r < R) \sim -a/r_0$, and

$$P_3(R) \sim \left(\frac{a}{R}\right)^6 \left(\frac{4\pi R^3 n}{3}\right)^2 = \left(\frac{4\pi na^3}{3}\right)^2, \quad (13)$$

At present it is unclear how the behavior can be matched on to the a^4 dependence predicted for $n|a|^3 \ll 1$ (see, e.g., Ref. [9]).

We next estimate the depletion of the condensate. In the LOCV approximation, the condensate fraction is given by [17]

$$\frac{n_0}{n} = 1 - n \int [1 - f(r)]^2 d^3r. \quad (14)$$

Substituting Eq. (7) in the above equation gives

$$\frac{n_0}{n} = \left(\frac{d}{r_0}\right)^3 \left[\frac{6}{\kappa^3} (\sin \kappa - \kappa \cos \kappa) - 1 \right]. \quad (15)$$

The condensate fraction calculated using the LOCV approximation is shown in Fig. 2 along with the predictions for the dilute limit given by [16]

$$\frac{n_0}{n} = 1 - \frac{4}{\sqrt{3}\pi} \left(\frac{a}{r_0}\right)^{3/2}. \quad (16)$$

Results for the hard-sphere potential obtained with the LOCV method, and exact quantum Monte Carlo calculations [12] are also shown in Fig. 2. The LOCV results for n_0/n are rather crude, since long-range correlations neglected in the LOCV approximation have a significant effect on n_0 . All the results indicate that superfluidity will be quenched when $a \gtrsim r_0$. In the case of hard-spheres it is known that the liquid solidifies at $r_0 \sim a$ [15].




FIG. 2. Condensate fraction vs. a/r_0 . LOCV calculations for attractive short range (full line) and hard-sphere (dashed line) interactions, Eq. 16 (dotted line) and exact QMC results for hard-spheres (stars).

The chemical potential at zero temperature is given by $\mu = dE/dN$. In the dilute limit the leading contributions to μ are $\mu = 4\pi\hbar^2 na/m [1 + (32/3)(na^3/\pi)^{1/2}]$, while, in the dense limit, the LOCV approximation gives

$$\mu = 22.22 \frac{\hbar^2 n^{2/3}}{m}. \quad (17)$$

The calculated $\mu(n)$ is shown in Fig. 3.

The square of the sound speed c_s is given by $c_s^2 = (n/m)d\mu/dn$, and this is also shown in Fig. 3. Its dilute and dense limits are given by $c_s^2 = 4\pi\hbar^2 na/m^2 [1 + 16(na^3/\pi)^{1/2}]$ and

$$c_s^2 = 14.81 \frac{\hbar^2 n^{2/3}}{m^2}, \quad (18)$$

respectively. At high densities the collective modes in the hydrodynamic limit are similar to those for fermions [18] since the energy per particle scales with density as $n^{2/3}$, as it does for a free Fermi gas. The energy per particle in a one-component Fermi gas, $E/N = (3/10)\hbar^2(6\pi^2n)^{2/3}/m$, is smaller by a factor 2.92 than the LOCV result in the dense Bose limit. The chemical potential and sound speed squared are therefore correspondingly larger in a dense Bose gas.




FIG. 3. The zero temperature chemical potential multiplied by $2mr_0^2/\hbar^2$ for: attractive short range (solid line) and hard-sphere (dashed line) from LOCV calculations, low density expansion (dotted line) and the square of hydrodynamic sound speed multiplied by $2m^2r_0^2/\hbar^2$ for: LOCV for attractive short range (double dash-dot), LOCV for hard-sphere (dash-double dot) and low density expansion (dash-dot).

The calculations of the energy density and the estimate of the three-body cluster probability $P_3(R)$ indicate that in limit where the scattering length is large compared to the interparticle spacing, the scattering length becomes irrelevant. These predictions could be investigated experimentally by measuring the size of a cloud of condensate and the particle lifetime as the scattering length is varied by exploiting a Feshbach resonance.

We thank G. Baym, S. Jonsell, and J. Yngvason for discussions. One of us, IEM, acknowledges financial support from the program “Universities of Russia” (project 015.01.01.04) and the Russian Ministry of Education grant. no. E00–3–12. The work of SC, JM, and VRP is supported in part by US National Science Foundation via grant PHY 98-00978.

- [1] S. Inouye, M. R. Andrews, J. Stenger, H.-J. Miesner, D. M. Stamper-Kurn, and W. Ketterle, *Nature* **392**, 151 (1998); Ph.Courteille, R. S. Freeland, and D. J. Heinzen, *Phys. Rev. Lett.* **81**, 69 (1998); J. L. Roberts, N. R. Claussen, J. P. Burke, C. A. Greene, E. A. Cornell, and C. A. Wieman, *Phys. Rev. Lett.* **81**, 5109 (1998). J. Stenger, S. Inouye, M. R. Andrews, H.-J. Miesner, D. M. Stamper-Kurn, W. Ketterle, *Phys. Rev. Lett.* **82**, 2422 (1999). J. L. Roberts, N. R. Claussen, S. L. Cornish, E. A. Donley, E. A. Cornell, and C. A. Wieman, *Phys. Rev. Lett.* **86**, 4211 (2001).
- [2] S. L. Cornish, J. L. Roberts, N. R. Claussen, J. L. Roberts, E. A. Cornell, and C. A. Wieman, *Phys. Rev. Lett.* **85**, 1795 (2000);
- [3] H. Heiselberg, *Phys. Rev. A* **63**, 043606 (2001).
- [4] W. Lenz, *Z. Phys.* **56**, 778 (1929).
- [5] T. D. Lee and C. N. Yang, *Phys. Rev.* **105**, 1119 (1957). T. D. Lee, K. Huang and C. N. Yang, *Phys. Rev.* **106**, 1135 (1957).
- [6] T. T. Wu, *Phys. Rev.* **115**, 1390 (1959).
- [7] L. P. Pitaevskii and S. Stringari, *Phys. Rev. Lett.* **81**, 4541 (1998); *Phys. Rev. Lett.* **83**, 4237 (1999).
- [8] E. Braaten and A. Nieto, *Phys. Rev. B* **55**, 8090 (1997); E. Braaten, H.-W. Hammer, and S. Hermans, *cond-mat/0012043*.
- [9] P. F. Bedaque, E. Braaten, and H.-W. Hammer, *Phys. Rev. Lett.* **85**, 908 (2000); E. Braaten, and H.-W. Hammer, *cond-mat/0103331*
- [10] R. Jastrow, *Phys. Rev.* **98**, 1479 (1955).
- [11] M. H. Kalos, D. Levesque and L. Verlet, *Phys. Rev. A* **9**, 2178 (1974).
- [12] S. Giorgini, J. Boronat and J. Casulleras, *Phys. Rev. A* **60**, 5129 (1999).
- [13] V. R. Pandharipande, *Nucl. Phys. A* **174**, 641 (1971); V. R. Pandharipande, *Nucl. Phys. A* **178**, 123 (1971). V.R. Pandharipande and H.A. Bethe, *Phys. Rev. C* **7**, 1312 (1973).
- [14] V. R. Pandharipande and K. E. Schmidt, *Phys. Rev. A* **15**, 2486 (1977).
- [15] J-P Hansen, D. Levesque and D. Schiff, *Phys. Rev. A* **3**, 776 (1971).
- [16] S. T. Beliaev, *JETP* **34**, 433 (1958).
- [17] M. L. Ristig, P. M. Lam, and J. W. Clark, *Phys. Lett.* **55A**, 101-103 (1975).
- [18] G. M. Bruun and K. Burnett, *Phys. Rev. A* **58**, 2427 (1998).